

AD-A261 014



DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

(2)

It is estimated to average 1 hour per response including the time for reviewing instructions, searching existing data sources, gathering and reviewing the collection of information, sending comments regarding this burden estimate or any other aspect of this burdening this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Avenue, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

2. REPORT DATE

Oct. 12, 1992

3. REPORT TYPE AND DATES COVERED

Final

4. TITLE AND SUBTITLE

Time Series Analysis and Multivariate Analysis

5. FUNDING NUMBERS

DAA03-89-K-0033

6. AUTHOR(S)

Theodore W. Anderson

DTIC

ELECTE

FEB 22 1993

S C D

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Stanford University
Stanford, CA 943058. PERFORMING ORGANIZATION
REPORT NUMBER

9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park, NC 27709-221110. SPONSORING/MONITORING
AGENCY REPORT NUMBER

ARO 26394.8-MA

11. SUPPLEMENTARY NOTES

The view, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.

12a. DISTRIBUTION/AVAILABILITY STATEMENT

Approved for public release; distribution unlimited.

12b. DISTRIBUTION CODE

13. ABSTRACT (Maximum 200 words)

Final Report: Summary of results, list of scientific personnel; list of technical reports, and list of publications.

93-03282



14. SUBJECT TERMS

Asymptotic distributions, autocorrelations, goodness-of-fit tests, spectral distributions, elliptically contoured distributions.

15. NUMBER OF PAGES

4 pages

16. PRICE CODE

17. SECURITY CLASSIFICATION
OF REPORT

UNCLASSIFIED

18. SECURITY CLASSIFICATION

UNCLASSIFIED

19. SECURITY CLASSIFICATION
OF ABSTRACT

UNCLASSIFIED

20. LIMITATION OF ABSTRACT

UL

932818163

TIME SERIES ANALYSIS AND MULTIVARIATE ANALYSIS

FINAL REPORT

Theodore W. Anderson
Principal Investigator

Jan. 15, 1989, to July 14, 1992

U. S. Army Research Office
Contract DAAG29-89-K-0033

DEPARTMENT OF STATISTICS
STANFORD UNIVERSITY
STANFORD, CALIFORNIA

DTIC QUALITY INSPECTED 3

Accession For	
NTIS CRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input checked="checked" type="checkbox"/>
Unannounced	<input checked="checked" type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

THE VIEW, OPINIONS, AND/OR FINDINGS CONTAINED IN THIS REPORT ARE THOSE OF THE AUTHORS(S) AND SHOULD NOT BE CONSTRUED AS AN OFFICIAL DEPARTMENT OF THE ARMY POSITION, POLICY, OR DECISION, UNLESS SO DESIGNATED BY OTHER DOCUMENTATION.

Research was carried out mainly in areas of time series analysis and multivariate analysis. The most important results in the first area apply to regression, autoregression, moving averages, and more generally, stationary stochastic processes. In the second area the results have to do with elliptically contoured distributions.

A simple condition has been derived for a numerical-valued Markov chain to be first-order autoregressive. The stationary probabilities of a general finite chain can be efficiently estimated by the mean of the corresponding vector-valued process.

Statistical inference about regression models and autoregression models is often based on the estimators of the coefficients being approximately normally distributed – at least in large samples. The applicability of inference based on asymptotic theory has been substantially extended; a Lindeberg-type condition is sufficient to justify this. The disturbance vectors need not have a common covariance matrix nor do they need to be independent; they can be martingale increments. The condition is implied by a strongly uniformly integrable condition.

A very general asymptotic distribution of a (finite) set of autocorrelations has been obtained based on the theory for autoregression coefficients. The results for autocorrelations are valid for linear processes with martingale difference disturbances.

Goodness-of-fit tests for time series models based on standardized spectral distributions have been developed for testing patterns of dependence in stationary processes. Tests can be conducted on the basis of the difference between the sample standardized spectral distribution and hypothesized process standardized spectral distribution. Considered as a stochastic process with frequency as the “time” parameter, this difference multiplied by the square root of the sample size converges weakly to a Gaussian stochastic procession on $[0, \pi]$ with expected value 0. Because the process depends on autocorrelations, the conditions for the limit results are very weak. For a linear process (that is, $y_t = \sum_{s=0}^{\infty} \gamma_s r_{t-s}$) Anderson has shown the limiting distribution of the autocorrelations for the disturbances being martingale differences.

The Gaussian stochastic process is characterized by its covariance function. After a suitable monotonic transformation of the frequency parameter, the covariance function of the transformed process is a constant times $\min(u, v) - uv + q(u)q(v)$, $0 \leq u, v \leq 1$, where $q(u)$ depends on the hypothetical spectral distribution. Note that

$\min(u, v) - uv$ is the covariance function of the Brownian bridge.

Tests are based on functionals of the process, such as the Cramér-von Mises statistic, the Kolmogorov-Smirnov statistic, and the Anderson-Darling statistic. The distribution of the Cramér-von Mises or Anderson-Darling statistic depends on the eigenvalues of an integral equation involving the covariance function of the limiting process; the integral equation can be converted to a second-order differential equation. A formal solution to this problem for standardized spectral distributions has been found, although it is feasible only in certain cases. The distribution of the Kolmogorov-Smirnov criterion satisfies a partial differential equation; a numerical solution is provided by solving recursively an approximating difference equation.

This asymptotic theory has been justified under very general conditions because the distributions of the criteria depend on the asymptotic distributions of autocorrelations.

The principal investigator with Raül P. Mentz studied iterative methods for estimating the parameters of a Gaussian moving average process of order 1. These procedures yield the exact maximum likelihood estimates. Efficient calculation is based on the calculation of quadratic forms of relevant matrices in the time domain and the traces of related matrices in the frequency domain. In iterative procedures for exact maximum likelihood estimation in first-order Gaussian moving average model "Woodbury's formula" gives an efficient method for evaluating quadratic forms in the inverse of a covariance matrix. The computation of the trace of the inverse of a covariance matrix can be expressed as an infinite series in the autocorrelation. A very accurate approximation has been obtained.

The other topic of research is inference in elliptically contoured distributions. It was shown that a scale-invariant vector-valued function has the same asymptotic distribution as in the normal case except for a factor depending on the kurtosis of the distribution; this factor can be consistently estimated. A scale-invariant scalar-valued function with value and first derivatives 0 at the true parameter value has the same asymptotic distribution as in the normal case except for the kurtosis factor. In most cases this asymptotic distribution is χ^2 .

Scientific Personnel

T. W. Anderson, Principal Investigator
Rebecca Betensky
Enapadam Venkatraman
Douglas Haney
Qizhi Wang

Technical Reports*

22. Second-Order Moments of a Stationary Markov Chain and Some Applications, by T. W. Anderson, February 1989.
23. Asymptotic robustness in Regression and Autoregression Based on Lindeberg Conditions, by T. W. Anderson and Naoto Kunitomo, June 1989.
24. Theory and Applications of Elliptically Contoured and Related Distributions, by T. W. Anderson and Kai-Tai Fang, September 1990.
25. Iterative Procedures for Exact Maximum Likelihood Estimation in the First-Order Gaussian Moving Average Model, by T. W. Anderson and R. P. Mentz, November 1990.
26. The Asymptotic Distributions of Autoregressive Coefficients, by T. W. Anderson, April 1991.
27. Goodness of Fit Tests for Spectral Distributions, by T. W. Anderson, November 1991.
28. Nonnormal Multivariate Distributions: Inference Based on Elliptically Contoured distributions, by T. W. Anderson, July 1992.

Publications

Second-order moments of a Markov chain and some applications, by T. W. Anderson, *Probability, Statistics, and Mathematics: Papers in Honor of Samuel Karlin* (T. W. Anderson, Krishna B. Athreya, and Donald L. Iglehart, eds.), Academic Press, 1989, 1-16. [Technical Report No. 22]

*Technical reports Nos. 1-21 were issued on Contracts DAAG29-82-K-0156 and DAAG29-85-K-0239.

Inference in multivariate elliptically contoured distributions based on maximum likelihood, by T. W. Anderson and Kai-Tai Fang, *Statistical Inference in Elliptically Contoured and Related Distributions* (Kai-Tai Fang and T. W. Anderson, eds.), Allerton Press, Inc., New York, 1990, 201-216. [Technical Report No. 1]

Noncentral distributions of quadratic forms for elliptically contoured distributions, by H. Hsu, *Statistical Inference in Elliptically Contoured and Related Distributions* (Kai-Tai Fang and T. W. Anderson, eds.), Allerton Press, Inc., New York, 1990, 97-102. [A part of Technical Report No. 14]

Generalized T^2 -test for multivariate elliptically contoured distributions, by H. Hsu, *Statistical Inference in Elliptically Contoured and Related Distributions* (Kai-Tai Fang and T. W. Anderson, eds.), Allerton Press, Inc., New York, 1990, 243-255. [A part of Technical Report No. 14]

Invariant tests for multivariate elliptically contoured distributions, by H. Hsu, *Statistical Inference in Elliptically Contoured and Related Distributions* (Kai-Tai Fang and T. W. Anderson, eds.), Allerton Press, Inc., New York, 1990, 257-274. [A part of Technical No. 14]

Relationships among classes of spherical matrix distributions, by K. T. Fang and H. F. Chen, *Statistical Inference in Elliptically Contoured and Related Distributions* (Kai-Tai Fang and T. W. Anderson, eds.), Allerton Press, Inc., New York, 1990, 25-36. [Technical Report No. 10]

The asymptotic distribution of autocorrelation coefficients, by T. W. Anderson, *The Art of Statistical Science: a Tribute to G. S. Watson* (K. V. Mardia, ed.), John Wiley & Sons, Ltd., Chichester, Sussex, 1992, 9-25. [Technical Report No. 26]

Theory and applications of elliptically contoured and related distributions, by T. W. Anderson and K. T. Fang, *The Development of Statistics: Recent Contributions from China* (X. R. Chen, K. T. Fang and C. C. Yang, eds.), Longman Scientific and Technical, Harlow, Essex, 1992, 41-62. [Technical Report No. 24]

Asymptotic distributions of regression and autoregression coefficients with martingale difference disturbances, by T. W. Anderson and Naoto Kunitomo, *Journal of Multivariate Analysis*, 40 (1992), 221-243. [Technical Report No. 23]